



Evidence in Motion, LLC
13201 Magisterial Drive
Louisville, KY 40223 (USA)
Ph: (888) 709-7096 Fax: (502) 244-6813
info@evidenceinmotion.com
www.evidenceinmotion.com
Blog with us at blog.evidenceinmotion.com!

The logo for the Foundation for Physical Therapy, featuring a stylized blue and white graphic of a person's head and neck in profile, with a blue swoosh above it. To the right of the graphic, the text "Foundation for Physical Therapy" is written in a serif font. Below this, in a smaller sans-serif font, is the text "Percentage of profits donated to Foundation for Physical Therapy".
Foundation for
Physical Therapy
Percentage of profits
donated to Foundation for
Physical Therapy

Lower Extremity Reference List

Hip:

- Altman R, Alarcon G, Appelrouth D et al. The American College of Rheumatology criteria for the classification and reporting of osteoarthritis of the hip. *Arthritis and Rheumatism* 2003;34:505-14.
- Bierma-Zeinstra S, Bohnen A, Ramlal R et al. Comparison between two devices for measuring hip joint motions. *Clinical Rehabilitation* 1998;12:497-505.
- Biji D, Dekker J, van Baar M et al. Validity of Cyriax's concept *capsular pattern* for the diagnosis of osteoarthritis of hip and/or knee. *Scand J Rheumatol* 1998;27:347-51.
- Birrell F, Croft P, Cooper C et al. Prediction radiographic hip osteoarthritis from range of movement. *Rheumatology* 2001;40:506-12.
- Browder D, Enseki K, Fritz J. Intertester reliability of hip range of motion measurements and special tests. *J Orthop Sports Phys Ther.* 34(1), A1. 2004.
- Bullock-Saxton J, Bullock M. Repeatability of muscle length measures around the hip. *Physiotherapy Canada* 1994;46:105-9.
- Cliborne AV, Wainner RS, Rhon DI et al. Clinical hip tests and a functional squat test in patients with knee osteoarthritis: Reliability, prevalence of positive test findings, and short-term response to hip mobilization. *J Orthop Sports Phys Ther* 2004;34:676-85.
- Fishman L, Dombi G, Michaelson C et al. Piriformis syndrome: Diagnosis, treatment and outcome- a 10-year study. *Arch Phys Med Rehabil* 2002;83:295-301.
- Fishman L, Zybert P. Electrophysiologic evidence of piriformis syndrome. *Arch Phys Med Rehabil* 1992;73:359-64.
- Hoeksma HL, Dekker J, Ronday HK, et al. A Comparison of Manual Therapy and Exercise Therapy in Osteoarthritis of the Hip: A Randomized Clinical Trial. *Arthritis Rheum.* 2004;51:722-729.
- Hoeksma HL, Dekker J, Ronday HK, et al. Manual Therapy in Osteoarthritis of the Hip: Outcomes in sub-groups of patients. *Rheum.* 2005;44:461-4.
- Jari S, Paton R, Srinivasan M. Unilateral limitation of abduction of the hip: A valuable clinical sign for DDH? *J Bone Joint Surg* 2002;84-B:104-7.
- Joe G, Kovacs J, Miller K et al. Diagnosis of avascular necrosis of the hip in asymptomatic HIV-infected patients: Clinical correlation of physical examination with magnetic resonance imaging. *Journal of Back and Musculoskeletal Rehabilitation* 2002;16:135-9.

Klassabo M, Harms-Ringdahl K, Larsson G. Examination of passive ROM and capsular patterns in the hip. *Physiotherapy Research International* 2003;8:1-12.

Lin Y-C, Davey R, Cochrane T. Tests for physical function of the elderly with knee and hip osteoarthritis. *Scand J Med Sci Sports* 2001;11:280-6.

Loudon JK. Case Report: Manual therapy management of hip osteoarthritis. *J Man Manip Ther.* 1999;7:203-208.

MacDonald CW, Whitman JW, Cleland JM, Smith MB, Hoeksma HL. Clinical Outcomes Following Manual Physical Therapy and Exercise for Hip Osteoarthritis. A Case Series. *J Orthop Sports Phys Ther.* 2006; 36:588-599.

Roddy E, Zhang W, Doherty M, et al. Evidence-based recommendations for the role of exercise in the management of osteoarthritis of the hip or knee – the MOVE consensus. *Rheumatology.* 2005;44,67-73.

Whitman JM, Flynn TW, Childs JD, Wainner RS, Gill HE, Ryder MG, Garber MB, Bennett AC, Fritz JM. A Comparison Between Two Physical Therapy Treatment Programs for Patients with Lumbar Spinal Stenosis: A Randomized Clinical Trial. *Spine.* In Press.

Knee:

Reference List

Altman RD, Bloch DA, Bole GG, Jr., et al. Development of clinical criteria for osteoarthritis. *J Rheumatol.* 1987;14 Spec No:3-6.

Bachman L, Haberzeth S, Stuerer J, ter Riet G. The accuracy of the Ottawa knee rule to rule out knee fractures: A systematic review. *Ann Intern Med.* 2004;140:121-4.

Baker KR, Nelson ME, Felson DT, Layne JE, Sarno R, Roubenoff R. The efficacy of home based progressive strength training in older adults with knee osteoarthritis: a randomized controlled trial. *J Rheumatol.* 2001;28(7):1655-65.

Bellamy N, Buchanan WW, Goldsmith CH, Campbell J, Stitt LW. Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *J Rheumatol.* 1988;15(12):1833-40.

Boeree N, Ackroyd C. Assessment of the menisci and cruciate ligaments: an audit of clinical practice. *Injury.* 1991;22:291-4.

Bonamo J, Shulman G. Double contrast arthrography of the knee. A comparison to clinical diagnosis and arthroscopic finding. *Arthrography and Arthroscopy.* 1988;11:1041-6.

Braunstein E. Anterior cruciate ligament injuries: a comparison of arthrographic and physical diagnosis. *AJR Am J Roentgenol.* 1982;138:423-5.

Bulloch B, Neto G, Plint A, Lim R, Lidman P, Reed M, et al. Validation of the Ottawa Knee Rule in children: A multicenter study. *Ann Emerg Med.* 2003;42:48-55.

Cleland JA, McRae M. Patellofemoral pain syndrome: A critical analysis of current concepts. *Physical Therapy Reviews.* 2002;7:153-61.

- Chang RW, Falconer J, Stulberg SD, Arnold WJ, Manheim LM, Dyer AR. A randomized, controlled trial of arthroscopic surgery versus closed-needle joint lavage for patients with osteoarthritis of the knee. *Arthritis Rheum.* 1993;36(3):289-96.
- Cook J, Khan K, Kiss Z, Puram C, Griffiths L. Reproducibility and clinical utility of tendon palpation to detect patellar tendinopathy in young basketball players. *British Journal of Sports Medicine.* 2001;35:65-72.
- Dervin G, Stiell I, Wells G, Rody K, Grabowski J. Physicians' accuracy and interrater reliability for the diagnosis of unstable meniscal tears in patients having osteoarthritis of the knee. *Canadian Medical Association.* 2001;44:267-74.
- Deyle GD, Allison SC, Ryder MG, Matekel RL, Stang JM, Gohdes DD, Hutton JP, Henderson NE, Garber MB. Physical Therapy treatment effectiveness for osteoarthritis of the knee: a comparison of supervised clinical exercise and manual therapy procedures versus a home exercise program. *Phys Ther.* 2005; 85(12):1301-1317.
- Deyle GD, Henderson NE, Matekel RL, Ryder MG, Garber MB, Allison SA. Effectiveness of Manual Physical Therapy and Exercise in Osteoarthritis of the Knee: A Randomized, Controlled Trial. *Ann Int Med.* 2000; 132(3): 173-181.
- Emparanza J, Aginaga J. Validation of the Ottawa Knee Rules. *Ann Emerg Med.* 2001;38:364-8.
- Eren O. The accuracy of joint line tenderness by physical examination in the diagnosis of meniscal tears. *Arthroscopy.* 2003;19:850-4.
- Ettinger WH, Jr., Burns R, Messier SP, et al. A randomized trial comparing aerobic exercise and resistance exercise with a health education program in older adults with knee osteoarthritis. The Fitness Arthritis and Seniors Trial (FAST). *JAMA* 1997;277(1):25-31.
- Evans P, Bell D. Prospective evaluation of the McMurray test. *Am J Sports Med.* 1993;21:604-9.
- Fitzgerald G, McClure P. Reliability of measurements obtained with four tests for patellofemoral alignment. *Phys Ther.* 1995;75:84-92.
- Fowler P, Lubliner J. The predictive value of 5 clinical signs in the evaluation of meniscal pathology. *Arthroscopy.* 1989;5:184-6.
- Fransen M, McConnell S, Bell M. Therapeutic exercise for people with osteoarthritis of the hip or knee. A systematic review. *J Rheumatol.* 2002;29(8):1737-45.
- Fritz JM, Delitto A, Erhard RE, Roman M. An examination of the selective tissue tension scheme, with evidence for the concept of a capsular of the knee. *Physical Therapy.* 1998;78:1046-61.
- Garvin G, Munk P, Vellet A. Tears of the medial collateral ligament: magnetic resonance imaging findings and associated injuries. *Can Assoc Radiol J.* 1993;44:199-204.
- Hayes K, Petersen C. Reliability of assessing end-feel and pain and resistance sequence in subjects with painful shoulders and knees. *J Orthop Sports Phys Ther.* 2001;31:432-45.
- Hayes K, Petersen C, Falconer J. An examination of Cyriax's passive motion tests with patients having osteoarthritis of the knee. *Phys Ther.* 1994;74:697-709.
- Jones A, Hopkinson N, Patrick M, Berman P, Doherty M. Evaluation of a method for clinically assessing osteoarthritis of the knee. *Annals of the Rheumatic Diseases.* 1992;51:243-5.

- Katz J, Fingerhuth R. The diagnostic accuracy of ruptures of the anterior cruciate ligament comparing the Lachman test, the anterior drawer sign, and the pivot shift test in acute and chronic knee injuries. *Am J Sports Med.* 1986;14:88-91.
- Ketelslegers E, Collard X, Vande Berg B, et al. Validation of the Ottawa Knee Rules in an emergency teaching center. *Eur Radiol.* 2002;12:1218-20.
- Khine H, Dorfman D, Avner J. Applicability of Ottawa knee rule for knee injury in children. *Pediatric Emergency Care.* 2001;17:401-4.
- Kim S, Kim H. Reliability of the anterior drawer test, the pivot shift test, and the Lachman test. *Clinical Orthopaedics and Related Research.* 1995;317:237-42.
- Lee J, Yao L, Phelps C, Wirth C, Czajka J, Lozman J. Anterior cruciate ligament tears: MR imaging compared with arthroscopy and clinical tests. *Radiology.* 1988;166:861-4.
- Liu S, Osti L, Henry M, Bocchi L. The diagnosis of acute complete tears of the anterior cruciate ligament. *J Bone Joint Surg.* 1995;77:586-8.
- Muellner T, Weinstabl R, Schabus R, Vecsei V, Kainberger F. The diagnosis of meniscal tears in athletes. A comparison of clinical and magnetic resonance imaging investigations. *Am J Sports Med.* 1997;25:7-12.
- Puett DW, Griffin MR. Published trials of nonmedicinal and noninvasive therapies for hip and knee osteoarthritis. *Ann Intern Med.* 1994;121(2):133-40.
- Rubinstein R, Shelbourne D. The accuracy of the clinical examination in the setting of posterior cruciate ligament injuries. *Am J Sports Med.* 1994;4.
- Shelbourne D, Martini D. Correlation of jointline tenderness and meniscal lesions in patients with acute anterior cruciate ligament tears. *Am J Sports Med.* 1995;23:166-70.
- Stiell I, Greenberg G, Wells G, et al. Derivation of a decision rule for the use of radiography in acute knee injuries. *Ann Emerg Med.* 1995;26:405-13.
- Stiell I, Wells G, Hoag R, Sivilotti M, Cacciotti T, Verbeek R, et al. Implementation of the Ottawa Knee Rule for the use of radiography in acute knee injuries. *JAMA.* 1997;278:2075-9.
- Sutlive T, Mitchell S, Maxfield S, McLean C, Neumann J, et al. Identification of individuals with patellofemoral pain whose symptoms improved after a combined program of foot orthosis use and modified activity: A preliminary investigation. *Phys Ther.* 2004;84:49-61.
- Watson C, Propps M, Galt W, Redding A, Dobbs D. Reliability of McConnell's classification of patellar orientation in symptomatic and asymptomatic subjects. *J Orthop Sports Phys Ther.* 1999;29:378-93.
- van Baar ME, Assendelft WJ, Dekker J, Oostendorp RA, Bijlsma JW. Effectiveness of exercise therapy in patients with osteoarthritis of the hip or knee: a systematic review of randomized clinical trials. *Arthritis Rheum.* 1999;42(7):1361-9.

Foot and Ankle:

Reference List

- Alonso A, Khoury L, Adams R. Clinical tests for ankle syndesmosis injury: Reliability and prediction of return to function. *J Orthop Sports Phys Ther.* 1998;27:276-84.
- Bachman L, Kolb E, Koller M, Steurer J, ter Riet G. Accuracy of Ottawa ankle rules to exclude fractures of the ankle and mid-foot: systematic review. *BMJ.* 2003;326.
- Bennell K, Talbot R, Wajswelner H, Techovanich W, Kelly D. Intra-rater and inter-rater reliability of a weight-bearing lunge measure of ankle dorsiflexion. *Australian Physiotherapy.* 1998;44:175-80.
- Bennett JE, Reinking MF, Pluemer B, Pentel A, Seaton M, Killian C. Factors Contributing to the Development of Medial Tibial Stress Syndrome in High School Runners. *J Orthop Sports Phys Ther.* 2001;31:504-10.
- Beumer A, Swierstra B, Mulder P. Clinical diagnosis of syndesmotic ankle instability. Evaluation of stress tests behind the curtain. *Acta Orthop Scand.* 2002;73:667-9.
- Coetzer D, Brantingham J, Nook B. The Relative Effectiveness of Piroxicam Compared to Manipulation in the Treatment of Acute Grades 1 and 2 Inversion Ankle Sprains. *Journal of the Neuromusculoskeletal System.* 2001;9(1):1-12
- Dananberg HJ, Shearstone J, Guillano M. Manipulation method for the treatment of ankle equinus. *J Am Podiatr Med Assoc.* 2000;90(8):385-9.
- Dananberg HJ. Manipulation of the ankle as a method of treatment for ankle and foot pain. *J Am Podiatr Med Assoc.* 2004 Jul-Aug;94(4):395-9.
- De Garceau D, Dean D, Requejo S, Thorardson D. The association between diagnosis of plantar fasciitis and windlass test results. *Foot and Ankle International.* 2003;24:251-5.
- Eiff MP, Smith AT, Smith GE. Early mobilization versus immobilization in the treatment of lateral ankle sprains. *Am J Sports Med.* 1994; 22:83-88.
- Eisenhart AW, Gaeta TJ, Yens DP. Osteopathic manipulative treatment in the emergency department for patients with acute ankle injuries. *J Am Osteopath Assoc.* 2003 Sep;103(9):417-21.
- Gaebler C, Kukla C, Breitenseher M, Nellas Z, Mittlboeck M, Trattinig S, Vecsei V. Diagnosis of lateral ankle ligament injuries. Comparison between talar tilt, MRI and operative findings in 112 athletes. *Acta Orthop Scand.* 1997;63:286-90.
- Green T, Refshauge K, Crosbie J, Adams R. A randomized controlled trial of a passive accessory joint mobilization on acute ankle inversion sprains. *Phys Ther.* 2001;81:984-994.
- Hertel J, Denegar CR, Monroe MM, Stokes WL. Talocrural and subtalar joint instability after lateral ankle sprain. *Medicine & Science in Sports & Exercise.* 1999;13:1501-8.
- Hunt GC. Injuries of peripheral nerves of the leg, foot and ankle: an often unrecognized consequences of ankle sprains. *The Foot.* 2003;13:14-8.
- Hunt GC. *Ankle sprain in a 14-year-old-girl.* Edinburgh: Butterworth Heinemann; 2004.

Hunt GC, Sneed T, Hamann H, Chisam. Biomechanical and histological considerations for development of plantar fasciitis and evaluation of arch taping as a treatment option to control associated plantar heel pain: A single-subject design. *The Foot*. 2004.

Jennings J, Davies GJ. Treatment of cuboid syndrome secondary to lateral ankle sprains: a case series. *J Orthop Sports Phys Ther*. 2005;35(7):409-15.

Liu S, Nuccion S. Diagnosis of an anterolateral ankle impingement. *American Journal of Sports Medicine*. 1997;25:389-94.

Maffulli. The clinical diagnosis of subcutaneous tears of the Achilles tendon. *The American Journal of Sports Medicine*. 2004;26:266-70.

Menz H, Tiedemann A, Kwan M, Latt M, Sherrington C, Lord S. Reliability of clinical tests of foot and ankle characteristics in older people. *Journal of the American Podiatric Association*. 2003;93:380-7.

Payne C, Chuter V, Miller K. Sensitivity and specificity of the functional halluc limitus test to predict foot function. *Journal of the American Podiatric Association*. 2002;92:269-71.

Pellow JE, Brantingham JW. The efficacy of adjusting the ankle in the treatment of subacute and chronic grade I and grade II ankle inversion sprains. *J Manipulative Physiol Ther*. 2001; 24:17-24.

Requejo SM, Kulig K, Thordarson DB. Management of foot pain associated with accessory bones of the foot: two clinical case reports. *J Orthop Sports Phys Ther*. 2000;30(10):580-91;

Shamus J, Shamus E, Gugel RN, Brucker BS, Skaruppa C. The effect of sesamoid mobilization, flexor hallucis strengthening, and gait training on reducing pain and restoring function in individuals with hallux limitus: a clinical trial. *J Orthop Sports Phys Ther*. 2004;34(7):368-76.

Stiell I, Greenberg G, McKnight R, Nair R, McDowell I, et al. A study to develop clinical decision rules for the use of radiography in acute ankle injuries. *Ann Emerg Med*. 1992;21:384-90.

Stiell I, McKnight R, Greenberg G, Nair R, McDowell I, Wallace G. Interobserver agreement in the examination of acute ankle injury patients. *American Journal of Emergency Medicine*. 1992;10:14- Tropp H, Norlin R. Ankle performance after ankle fracture: a randomized study of early mobilization. *Foot Ankle Int*. 1995; 16:79-83.

van der Wees PJ, Lenssen AF, Hendriks EJ, Stomp DJ, Dekker J, de Bie RA. Effectiveness of exercise therapy and manual mobilisation in ankle sprain and functional instability: a systematic review. *Aust J Physiother*. 2006;52(1):27-37.

van Dijk C, Mol B, Lim L, Mati R, Bossuyt P. Diagnosis of a ligament rupture of the ankle joint. Physical examination, arthrography, stress radiography and sonography compared in 160 patients after inversion trauma. *Acta Orthop Scand*. 1996;67:566-70.

Van Gheluwe B, Kirby K, Roosen P, Phillips R. Reliability and accuracy of biomechanical measurement of the lower extremities. *Journal of the American Podiatric Association*. 2002;92:317-26.

Whitman JM, Childs JD, Walker V. The use of manipulation in a patient with an ankle sprain injury not responding to conventional management: a case report. *Man Ther*. 2005;10(3):224-31.

Young B, Walker MJ, Strunce J, Boyles R. A combined treatment approach emphasizing impairment-based manual physical therapy for plantar heel pain: a case series. *J Orthop Sports Phys Ther.* 2004;34(11):725-33.